arXiv:1112.3432v1 [math.CO] 15 Dec 2011

Line-transitive point-imprimitive linear spaces with Fang-Li parameter gcd(k, r) at most 10

Haiyan Guan, Delu Tian and Shenglin Zhou*
Department of Mathematics, South China University of Technology,
Guangzhou, Guangdong 510640, P. R. China

December 16, 2011

Abstract

This paper is a further contribution to the classification of line-transitive finite linear spaces. We prove that if $\mathcal{S}$ is a non-trivial finite linear space with the Fang-Li parameter $\gcd(k, r)$ is 9 or 10, the group $G \leq \mathrm{Aut}(\mathcal{S})$ is line-transitive and point-imprimitive, then $\mathcal{S}$ is the Desarguesian projective plane $\mathrm{PG}(2, 9)$.

Keywords: linear space, line-transitive, point-imprimitive

MR(2000) Subject Classification 05B05, 05B25, 20B25

1 Introduction

A finite linear space $\mathcal{S}$ is an incident structure $(\mathcal{P}, \mathcal{L})$ where $\mathcal{P}$ is a set of v points and $\mathcal{L}$ is a set of b distinguished subsets of $\mathcal{P}$ called lines, such that any two points are incident with exactly one line. A linear space $\mathcal{S}$ is said to be non-trivial if every line is incident with at least 3 points and $b \geq 2$.

An automorphism of $\mathcal{S}$ is a permutation of $\mathcal{P}$ which leaves the set $\mathcal{L}$ invariant. The full automorphism group of $\mathcal{S}$ is denoted by $\mathrm{Aut}(\mathcal{S})$. Any subgroup of $\mathrm{Aut}(\mathcal{S})$ is called an automorphism group of $\mathcal{S}$. If a subgroup G of $\mathrm{Aut}(\mathcal{S})$ acts transitively on the set of points (lines, flags), then we say that G is line-transitive (point-transitive, flag-transitive), where a flag is an incident point-line pair (α, λ). Similarly, the automorphism group G is said to be point-primitive (line-primitive) if it acts primitively on points (lines). If the sizes of all lines are equal, then we say that $\mathcal{S}$ is a regular linear space. In this paper, we assume that the automorphism group $G \leq \mathrm{Aut}(\mathcal{S})$ is line-transitive. It follows that $\mathcal{S}$ is regular, and the line size will be denoted by k. It is well-known that if G is line-transitive then it is also

---

[*] Partially supported by the NSFC(No:11071081). Corresponding author: slzhou@scut.edu.cn

point-transitive([3]), and for a non-trivial regular linear space every point lies on the same number r of lines.

In [2], the line-transitive point-imprimitive linear spaces with the Fang-Li parameter $\gcd(k,r) \leq 8$ are classified. As an application, C. E. Praeger and S. L. Zhou [20] give the classification of the line-transitive linear spaces with $k \leq 12$, extending the Camina-Mischke's one in [5].

This paper is a continuation of [2]. We extended the classification of line-transitive poin-imprimitive linear spaces with the Fang-Li parameter $\gcd(k,r)$ up to 10, and proved the following result.

Theorem 1.1 *Let $\mathcal{S}$ be a non-trivial finite linear space with v points and Fang-Li parameter $k^{(r)} = 9$ or 10. Assume that $G \leq \mathrm{Aut}(\mathcal{S})$ is line-transitive and point-imprimitive. Then $\mathcal{S}$ is the Desarguesian projective plane* $\mathrm{PG}(2,9)$.

The proof of Theorem 1.1 rely on the realize of the algorithms in [1] and the available theory of finite line-transitive, point-imprimitive linear spaces in [2, 19, 20].

Combining Theorem 1.1 and [2, Theorem 1.2], we get the following.

Corollary 1.2 *Suppose that $\mathcal{S}$ is a linear space admitting a line-transitive, point-imprimitive subgroup of automorphisms, and with the Fang-Li parameter $k^{(r)} \leq 10$. Then $\mathcal{S}$ is the Desarguesian projective plane* $\mathrm{PG}(2,4)$, $\mathrm{PG}(2,7)$ *or* $\mathrm{PG}(2,9)$, *or the Mills design or Colbourn-McCalla design, both with* $(v,k) = (91,6)$, *or one of the* 467 *Nickel-Niemeyer-O'Keefe-Penttila-Praeger designs with* $(v,k) = (729,8)$.

The structure of the paper is as follows. Section 2 contains some parameters and our hypotheses, Section 3 gives some elementary results on linear spaces which will be frequently used in the proof of our result. In Section 4, we used and realized the algorithms in [1] to produce a list of potential parameters and some group theoretic information for $k^{(r)} = 9$ or 10. We then made a detailed analysis of each of these potential parameters in Sections 5 and 6.

2 Definitions and Hypothesis

In this paper, we use the following definitions, notation and hypothesis.

2.1 Dlandtsheer-Doyen parameters

If G acts point-imprimitively on $\mathcal{P}$, then $\mathcal{P}$ has a non-trivial partition of imprimitivity. Let $\mathfrak{C} = \{C_1, C_2, \cdots, C_d\}$ be the partition where $|C_i| = c$ for all $1 \leq i \leq d$, so that $v = cd$. Then there exist two positive integers x, y, such that

$$c = \frac{\binom{k}{2} - x}{y} \quad \text{and} \quad d = \frac{\binom{k}{2} - y}{x},$$

where x is the number of inner pairs on a line ([11]). Here c, d, x, y are called the Dlandtsheer-Doyen parameters.

2.2 Fang-Li parameters

Let v, b, k be the parameters of $\mathcal{S} = (\mathcal{P}, \mathcal{L})$, and $r = \frac{v-1}{k-1}$ is the number of lines incident with a given point which is independent of the choice of point by the line-transitivity. Define

$$k^{(v)} = \gcd(k, v), k^{(r)} = \gcd(k, r), b^{(v)} = \gcd(b, v), b^{(r)} = \gcd(b, r).$$

Then

$$b = b^{(v)} b^{(r)}, k = k^{(v)} k^{(r)}, v = b^{(v)} k^{(v)}, r = k^{(r)} b^{(r)}.$$

Clearly, $k^{(r)} = k/\gcd(k, v) = \gcd(k, v-1)$. Here $k^{(v)}, k^{(r)}, b^{(v)}, b^{(r)}$ are called the Fang-Li parameters ([15]).

2.3 Top and Bottom groups

Let $C \in \mathfrak{C}$, $G^{\mathfrak{C}}$ be the transitive permutation group on $\mathfrak{C}$ induced by G, then $G^{\mathfrak{C}} = G/G_{(\mathfrak{C})}$ and it is called the top group([1]). Similarly, define the bottom group $G^C = G_C/G_{(C)}$ ([1]), which is the transitive group induced by G on $C \in \mathfrak{C}$. Moreover, G^C is independent of the choice of C in $\mathfrak{C}$ up to permutation isomorphism. For partitions $\mathfrak{C}, \mathfrak{C}'$ of $\mathcal{P}$, $\mathfrak{C}$ *refines* $\mathfrak{C}'$ if every class of $\mathfrak{C}$ is contained in a class of $\mathfrak{C}'$, and this refinement is strict if $\mathfrak{C} \neq \mathfrak{C}'$. We also say that $\mathfrak{C}'$ is *coarser or strictly coarser* than $\mathfrak{C}$. We call a partition minimal if it has no non-trivial strict refinement, maximal if the only G-invariant partition of $\mathcal{P}$ that is strictly coarser than $\mathfrak{C}$ is the trivial partition $\{\mathcal{P}\}$ with only one class. If a partition $\mathfrak{C}$ of $\mathcal{P}$ is both maximal and minimal, then we call G *2-step imprimitive* relative to $\mathfrak{C}$, in this case, both $G^{\mathfrak{C}}$ and G^C are primitive. So if G is 2-step imprimitive then $G^{\mathfrak{C}}$ and G^C are primitive groups of degree d and c respectively. Relying on the classification of primitive groups of degree less than 4096 (cf. [8, 13, 21]), if $c \leq 4096$ then we know the structure of the bottom group, similarly for d and the top group. This will be frequently used in Sections 5 and 6.

If the subgroup $G_{(\mathfrak{C})}$ is transitive on each class of $\mathfrak{C}$, then we say that the partition $\mathfrak{C}$ is G-normal. By [7], we know that if G is a line-transitive point-imprimitive group of automorphisms of a linear space, then either

(i) there exists a non-trivial G-normal point-partition; or

(ii) G is point-quasiprimitive and almost simple.

Recall that a finite permutation group G is *quasiprimitive* if every nontrivial normal subgroup of G is transitive. Equivalently, G is quasiprimitive if every minimal normal subgroup of G is transitive (since overgroups of transitive groups are transitive).

2.4 Intersection type and t_{max}

The intersection type and t_{max} will play an important role in our discussion. We give their definitions in the following.

Definition 2.1 ([10]) *Let $\mathcal{S}$ be a linear space, $\lambda \in \mathcal{L}$. Define*

$$d_i = |C \in \mathfrak{C} : |C \cap \lambda| = i|,$$

and

$$(0^{d_0}, 1^{d_1}, \cdots, k^{d_k})$$

to be the intersection type of $\mathcal{S}$.

We let the set of non-zero intersection sizes

$$Spec(\mathcal{S}) := \{i > 0 : d_i \neq 0\}.$$

is the spectrum of $\mathcal{S}$. Since the transitivity of G on $\mathcal{L}$, the intersection type and spectrum are independent of the choice of λ. We sometimes write $Spec_{\mathfrak{C}}(\mathcal{S})$ if we need to specify the partition $\mathfrak{C}$. Since $d_0 = d - \sum_{i=1}^{k} d_i$, then it does not matter to denote the intersection type of $\mathcal{S}$ by $(1^{d_1}, \cdots, k^{d_k})$.

Definition 2.2 ([1]) *For a given intersection type $(1^{d_1}, \cdots, k^{d_k})$, and non-empty subset $S \subseteq Spec(\mathcal{S})$, set $d(S) = \sum_{i \in \mathcal{S}} d_i$. Define t_{max} to be the largest positive integer t such that, for all $S \subseteq Spec(\mathcal{S})$, and all positive integers $h \leq min\{t, d(S)\}$,*

$$\prod_{j=0}^{h-1} (d - j) \mid b \prod_{j=0}^{h-1} (d(\mathcal{S}) - j).$$

2.5 Group theory notations and HYPOTHESIS

If $\mathfrak{C}$ is G-normal, then let $K = G_{(\mathfrak{C})}$, $S = \mathrm{Soc}(K)$, $X = C_G(K)$, and $Y = C_G(S)$ ([20]), where $\mathrm{Soc}(G)$ denotes the socle of the group G, that is, the product of the minimal normal subgroups of G. Throughout the paper we assume the following HYPOTHESES.

HYPOTHESIS

Let $\mathcal{S}$ be a non-trivial finite linear space with v points and b lines, each of size k, and with r lines through each point. Let $G \leq \mathrm{Aut}(\mathcal{S})$ be line-transitive and point-imprimitive such that $\mathfrak{C} = \{C_1, C_2, \cdots, C_d\}$ be a G-invariant partition of $\mathcal{P}$ with d classes of size c where $c > 1$ and $d > 1$. Also $\mathcal{S}$ has the Delandtsheer-Doyen parameters c, d, x, y, the Fang-Li parameters $k^{(v)}, k^{(r)}, b^{(v)}, b^{(r)}$, intersection type $(1^{d_1}, \cdots, k^{d_k})$, t_{max}, the subgroups K, S, X and Y if $\mathfrak{C}$ is G-normal defined as above.

3 Some Preliminary Results

The following lemma is essential for the proof of Theorem 1.1. So for the completeness, we give a detailed proof here.

Lemma 3.1 [5, Lemma 5] *Let G satisfy* HYPOTHESIS*, p be a prime.*

(i) If $p \mid |G_{(\lambda)}|$*, and* $k^2 - k + 1 > max(r + k - p + 1, r)$*, then* $p \mid |G^{\lambda}|$ *for any line* λ*.*

(ii) If $p > k$ *and* $k^2 - k + 1 > r$*, then* $p \mid v$ *or* $p \mid (v - 1)$*. Further if* T *is a Sylow* p*-subgroup of* G *then* $|T|$ *divides* v *or* $v - 1$ *respectively.*

Proof: (i) Let $p \mid |G_{(\lambda)}|$ and P be a Sylow p-subgroup of G_λ for $\lambda \in \mathcal{L}$. If $p \nmid |G^{\lambda}|$, then P fixes λ point-wise. So if P fixes two points, it has to fix the unique line point-wise containing the two points. If the set of fixed points $F = Fix_{\mathcal{P}}(P)$ is the line λ, by [6, Lemma 3], we conclude that G is flag-transitive, a contradiction. Thus F has a structure of regular linear space with line size k, therefore, $k^2 - k + 1 \leq |F|$. Furthermore, by [6, Lemma 1], $|F| \leq max(k + r - p - 1, r)$, a desired contradiction.
(ii) Let P be a p-subgroup of G. If $P \leq G_\lambda$ for $\lambda \in \mathcal{L}$. By (i) $p \mid |G^{\lambda}|$, but $G^{\lambda} \leq S_k$, this is impossible. Thus $P = 1$. Therefore , there is no p-subgroup fixing a line, then at most one point is fixed by P, that is to say $|T|$ divides v when it has no fixed point or $v - 1$ when it has one fixed point on $\mathcal{P}$. □

Note that if $p \mid |G_\lambda|$, (i) also holds.

For a primitive permutation group G, we call G is t-transitive if the integer t is the maximal integer such that G is t-transitive.

Lemma 3.2 [1, Lemma 4.7] *If* $G^{\mathfrak{C}}$ *is* t*-transitive then* $t \leq t_{max}$*.*

Lemma 3.3 [12, Proposition 2.6] *Assume that the* HYPOTHESIS *holds.*

(i) The number $b^{(r)}$ *divides each non-trivial subdegree of* $G^{\mathfrak{C}}$*, and in particular,* $rank(G^{\mathfrak{C}}) \leq 1 + \delta$*.*

(ii) The number $b^{(r)}$ *divides each non-trivial subdegree of* G^{C}*, and in particular,* $rank(G^{C}) \leq 1 + \gamma$*.*

(iii) Moreover, for $\alpha \in \mathcal{P}$*,* $b^{(r)}$ *divides each non-trivial subdegree of* $G^{\mathcal{P}}$ *and each orbit length of* G_α *in* $\lambda \in \mathcal{L}$ *and* $\alpha \in \lambda$*.*

Lemma 3.4 *Assume that the* HYPOTHESIS *holds.*

(i) [16, Proposition 3.6] *If* $\gamma = 2$*,* $b^{(r)}$ *is odd and* c *is not a prime power, then* G^{C} *is 2-transitive on* C*.*

(ii) [16, Proposition 3.7] *If* $\delta = 2$*,* $b^{(r)}$ *is odd and* d *is a prime power, then* $G^{\mathfrak{C}}$ *is not 2-transitive on* $\mathfrak{C}$*.*

Lemma 3.5 [6, Lemma 2] *Assume that* $\mathcal{S} = (\mathcal{P}, \mathcal{L})$ *is a non-trivial linear space admitting a line-transitive automorphism group* G*. Let* $\lambda \in \mathcal{L}$ *and* $H \leq G_\lambda$ *such that, for* $F := \mathrm{Fix}_{\mathcal{P}}(H)$*,*

(i) $2 \leq |F \cap \lambda| < |F|$*, and*

(ii) if $K \leq G_\lambda$ *and* $|\mathrm{Fix}_{\mathcal{P}}(K) \cap \lambda| \geq 2$*, and* H *and* K *are conjugate in* G*, then* H *and* K *are conjugate in* G_λ*.*

Then the deduced linear space $\mathcal{S}|_F = (F, \mathcal{L}|_F)$ has constant line size and $N_G(H)$ acts line-transitively on $\mathcal{S}|_F$, where $\mathcal{L}|_F = \{\lambda \cap F : \lambda \in \mathcal{L}, |\lambda \cap F| \geq 2\}$.

Lemma 3.6 [1, Corollary 4.4] *Assume that the* HYPOTHESIS *holds. Let $\lambda \in \mathcal{L}$ and p be a prime dividing $|G_\lambda|$. Let P be a Sylow p-subgroup of G_λ and $F = \mathrm{Fix}_{\mathcal{P}}(P)$. Suppose that $2 \leq |F \cap \lambda| < |F|$. Then*

(i) $N_G(P)$ is line-transitive on $\mathcal{D}|_F$.

(ii) $C|_F = \{C \cap F : C \in \mathfrak{C}, C \cap F \neq 0\}$ is an $N_G(P)$-invariant partition of F.

(iii) $|F| = f \cdot |C \cap F|$ where $f = |\mathfrak{C}|_F|$, $C \cap F \in \mathfrak{C}|_F$, and $|C \cap F| \geq 3$.

Lemma 3.7 [1, Theorem 6.1] *Assume that the* HYPOTHESIS *holds and that C is G-normal.*

(i) If $k > 2x + \frac{3}{2} + \sqrt{4x - \frac{7}{4}}$, then $G_{(\mathfrak{C})}$ is semiregular on points and lines, $|G_{(\mathfrak{C})}| = c$ is odd, and $d_1 > 0$.

(ii) If $x \leq 8$, then $G_{(\mathfrak{C})}$ has an abelian subgroup S of index at most 2 such that S is normal in G, semiregular on points, and $|S| = c$ is odd.

(iii) If either of the conditions of (i) or (ii) holds, and if $\mathfrak{C}$ is minimal, then c is an odd prime power and G^C is affine.

Lemma 3.8 [20, Lemma 2.2] *Assume that the* HYPOTHESIS *hold and that $\mathfrak{C}$ is G-normal and minimal.*

(a) Then $\mathfrak{C}$ is the set of S-orbits in $\mathcal{P}$ and

  (i) Either $Y \cap K = 1$, or S is elementary abelian and $Y \cap K = S$.

  (ii) Either $X \cap K = 1$, or S is elementary abelian. and $X \cap K = K = S$.

(b) Suppose in addition that $\mathfrak{C}$ is maximal, that $Y \cap K = S \neq Y$, and that one of the following conditions holds.

Condition	$\mathrm{Soc}(G^{\mathfrak{C}})$	*Extra Property*
1	*abelian*	$\gcd(c, d) = 1$
2	*non-abelian*	*Schur multiplier of a minimal normal subgroup of $G^{\mathfrak{C}}$ has no section isomorphic to S*

Then G has a normal subgroup $M = T \times S$ where T is a minimal normal subgroup of G and $T^{\mathfrak{C}}$ is minimal normal in $G^{\mathfrak{C}}$. Moreover either

  (i) T is non-abelian and transitive on $\mathcal{P}$, or

  (ii) the set $\mathfrak{C}'$ of T-orbits in $\mathcal{P}$ is a G-normal partition of $\mathcal{P}$ with $|\mathfrak{C}'| = c$ such that for $C \in \mathfrak{C}$ and $C' \in \mathfrak{C}'$, $|C'| = d$ and $|C \cap C'| = 1$. Moreover either M is regular on $\mathcal{P}$ or T is not semiregular on $\mathcal{P}$.

4 Computing parameters

Assuming the Hypothesis holds with $k^{(r)} = 9$ or 10, we get totally 304 potential parameter sets (called Lines)

$$(d, c, x, y, \gamma, \delta, k^{(v)}, k^{(r)}, b^{(v)}, b^{(r)}, \text{intersection type}, t_{\max}),$$

by applying Algorithms 1 and 2 in [1] which implemented in GAP([14]). For each Line, there is no other partition of $\mathcal{P}$ which is neither a strict refinement nor strict coarser than $\mathfrak{C}$, so the automorphism group is 2-step imprimitive relative to $\mathfrak{C}$.

In Section 5, we deal with 33 Lines with $k^{(r)} = 9$ listed in Table 1. In Section 6 the remaining 271 Lines which listed in Table 2.

In Table 1 and Table 2, we renumber these Lines, called Cases, according to the different parameters

$$(d, c, x, y, \gamma, \delta, k^{(v)}, k^{(r)}, b^{(v)}, b^{(r)})$$

other than the intersection type, so there are 11 Cases in Table 1 and 106 Cases in Table 2 needed to analysis.

It is worth mentioning that among the computation of the intersection types, we modified the algorithm by adding the restrictive condition $\frac{\alpha_i}{gcd(\alpha,\beta)} | d_i'$ of [1, Prop 4.5(vi)].

5 The case where $k^{(r)} = 10$

Proposition 5.1 *Let $\mathcal{S}$ be a non-trivial finite linear space with v points and Fang-Li parameter $k^{(r)} = 10$. Assume that $\mathcal{S}$ has a subgroup G of automorphism group which is line-transitive and point-imprimitive. Then $\mathcal{S}$ is the Desarguesian projective plane* PG(2, 9).

Proof. We deal with 11 Cases in Table 1 case by case.

Cases 1 and 3-5: In each of these Cases, $k = 10$. Recall that [20] classified the linear spaces with line size at most 12, so by [20, Theorem 1.1], if Case 1 or 4 holds, then $\mathcal{S}$ is the Desarguesian projective plane PG(2, 9), and Cases 3, 5 can not occur by [9].

Case 2: Assume Case 2 holds, then $\mathrm{Soc}(G^C)$ is $\mathrm{PSL}(2, 2^7)$, or A_{129}. It follows that there exists a prime $p = 127 \mid |G^C|$, and so $127 \mid |G|$, we also have $p > k = 30$, $k^2 - k + 1 > r$, however $p \nmid v(v-1) = 2^3 \cdot 3^3 \cdot 5 \cdot 29 \cdot 43$, contradicts Lemma 3.1. Similarly, Case 8 can not occur, because here $\mathrm{Soc}(G^{\mathfrak{C}})$ is $\mathrm{PSL}(2, 2^7)$, or A_{129}, and there exists a prime $p = 127$ satisfying the conditions of Lemma 3.1(ii) but $p \nmid v(v-1)$.

Cases 7, 10 and 11: Since $k^2 - k + 1 > r$, and there exists a prime $p > k$ which is listed in Table 1, such that $p \mid |G|$, but $p \nmid v(v-1)$, a contradiction. In Table 1, the symbol (T, p) or (B, p) denotes the contradiction by the Top or Bottom group and the prime p such that p divides the order of the top or bottom group but p does not divides $v(v-1)$.

Cases 6 and 9: **(a)** For each Case, the partition $\mathfrak{C}$ is G-normal, K is semiregular.

For Case 6, suppose first that G is point-quasiprimitive and almost simple, then $G^{\mathfrak{C}} \cong G$ is an almost simple primitive group of degree 49. As G is point-transitive additionally, we have that $v \mid |G|$. Hence $G \geq A_{49}$. But here $t_{max} = 1$, contradicting Lemma 3.2. So $\mathfrak{C}$ is G-normal relative to K. If K is not semiregular, as $d_1 \cdot d_2 = 14 \times 7$ in LINE 20 and $28 \times 7 > 0$ in LINE 21, by [19, Corollary 4.2] and [19, Lemma 5.1], we have

$$1 + \frac{r}{k} d_1(d_1 - 1) \leq |\mathrm{Fix}_{\mathcal{P}}(K_\alpha)| \leq d,$$

it follows that $d_1 \leq 1$. This is a contradiction because $d_1 = 14$ or 28 in this CASE. Thus K is semiregular, and $K \trianglelefteq G^C$, then K is the unique soluble minimal regular normal subgroup of primitive group G^C, so $K = Z_{13}^2$, and $Y \cap K = K = S, G/Y \leq \mathrm{Aut}(S) = \mathrm{GL}(2, 13)$.

For CASE 9, suppose that G is quasiprimitive and almost simple, then $G \cong G^{\mathfrak{C}}$ is an almost simple primitive group of degree 169. Since in CASE 9, $t_{max} = 2$, G is at most two transitive. However, checked by GAP([14]), we know that there is no such group satisfying the conditions. Thus the partition $\mathfrak{C}$ is G-normal relative to K. Moreover, in CASE 9, $k = 70$ and $x = 14$, so we have that

$$k > 2x + \frac{3}{2} + \sqrt{4x - \frac{7}{4}} = \frac{59 + \sqrt{217}}{2},$$

hence by Lemma 3.7, K is semiregular, and G is affine. Therefore $K = Z_7^2, K = S = Y \cap K = \mathrm{Soc}(G^C), G/Y \leq \mathrm{Aut}(S) = GL(2, 7)$.

(b) CASE 9 can not occur.

Suppose $Y^{\mathfrak{C}} = 1$, then $Y = Y \cap K = S, Y \leq K$, hence $|G^{\mathfrak{C}}| = |G/K| \mid |G/Y|$. It follows that $d = 169 \mid (7^2 - 1)(7^2 - 7)$, a contradiction. So, by [20, Lemma 2.1], there exists a normal subgroup M of G, such that $S < M \leq Y, M/S \cong M^{\mathfrak{C}}$ is a minimal subgroup of $G^{\mathfrak{C}}$, and $M = T \times S$.

(1) $\mathrm{Soc}(G^{\mathfrak{C}}) = Z_{13}^2$, then $M^{\mathfrak{C}} = Z_{13}^2$. In this case M is regular on point $\mathcal{P}$, with a unique Sylow 13-subgroup $T = Z_{13}^2$, and T is regular on $\mathfrak{C}$. Since the intersection type is $(1^{42}, 2^{14})$, all entries in the $\mathfrak{C}$-intersection type are at most 2, no element of S of order 7 fixes a line, for if $g \in S$, $o(g) = 7$ and $g \in G_\lambda$ for some $\lambda \in \mathcal{L}$, then g fixes $C \cap \lambda$ for any $C \in \mathfrak{C}$, particularly, g fixes $C \cap \lambda$ for which $|C \cap \lambda| = 1$, thus g has fixed points, contradicting the fact that S is semiregular. Similarly there is no element of order 13 in T fixing a line. Hence M is semirugular on $\mathcal{S}$. So $v \mid b$. However, in CASE 9, $\frac{b}{v} = \frac{b^{(r)}}{k^{(v)}} = \frac{12}{7}$, a contradiction.

(2) $\mathrm{Soc}(G^{\mathfrak{C}}) = A_{13}^2$ or $\mathrm{PSL}(3, 3)^2$. Here $\mathrm{Soc}(G^{\mathfrak{C}}) \neq A_{169}$ since $t_{max} = 2$. In this case $M^{\mathfrak{C}}$ is nonabelian.

(2.1) $\mathrm{Soc}(G^{\mathfrak{C}}) = A_{13}^2$. Then $\mathfrak{C} = \mathfrak{C}_1 \times \mathfrak{C}_2$, $\mathfrak{C} = \{C_{ij} \,|\, 1 \leq i \leq 13, 1 \leq j \leq 13\}$, $A_{13}^2 \leq G^{\mathfrak{C}} \leq S_{13} \wr S_2$ in product action. $\mathrm{Soc}(G^C)$ acts on $\mathfrak{C}$ as follow. Let $(h_1, h_2) \in A_{13}^2 \leq G^{\mathfrak{C}}$, and $C_{ij} \in \mathfrak{C}$. Then $C_{ij}^{(h_1,h_2)} = C_{i^{h_1} j^{h_2}}$. Let P be a Sylow 11-subgroup of G. Since $11 \nmid b$, $P \leq G_\lambda$ for some $\lambda \in \mathcal{L}$. Moreover, because $11 \nmid |GL(2, 7)|$, then if P fixes some C_{ij} setwise, it must fix C_{ij} piontwise. Hence $\mathrm{Fix}_{\mathcal{P}}(P)$ is a union of $\mathfrak{C}$-classes. Without loss generality, we assume that $P = \langle (1, 2, 3, \cdots, 11) \rangle \times \langle (1, 2, 3, \cdots, 11) \rangle$, so $\mathrm{Fix}_{\mathfrak{C}}(P) = \{C_{ij} | i, j = 12 \text{ or } 13\}$, thus $|\mathrm{Fix}_{\mathfrak{C}}(P)| = 4$. Furthermore, since the intersection type is $(1^{42}, 2^{14})$ and $P \leq G_\lambda$, P fixes setwise the 42 classes C such that $|C \cap \lambda| = 1$ and the 14 classes C setwise such that

$|C\cap\lambda| = 2$. Since $42 \equiv 9 \pmod{11}$ and $14 \equiv 3 \pmod{11}$, thus P must fix at least $9+3 = 12$ classes setwise, a contradiction.

(2.2) $\mathrm{Soc}(G^{\mathfrak{C}}) = PSL(3,3)^2$ and $M^{\mathfrak{C}} \cong T^{\mathfrak{C}} = T$. For the Schur Multiplier of PSL(3, 3) is trivial, by Lemma 3.8, either T is transitive on points or the set of T-orbits forms a G-normal partition $\mathfrak{C}'$ satisfying CASE 6. In the latter case, T must be abelian, a contradiction. Thus T is transitive on $\mathcal{P}$, then $c \mid |T|$, anther contradiction. Therefore CASE 9 can not occur. In Table 1 the symbol KSXY refers to the situation that the CASE is ruled out by Lemma 3.8.

(c) CASE 6 can not occur.

Suppose $Y^{\mathfrak{C}} = 1$, then $Y = Y \cap K = S, Y \le K$, hence $|G^{\mathfrak{C}}| = |G/K| \mid |G/Y|$ and so $|G^{\mathfrak{C}}| \mid |GL(2,7)|$. Therefore $d = 49 \mid (13^2-1)(13^2-13)$, a contradiction. By Lemma 3.8, there exists a normal subgroup M of G, such that $S < M \le Y, M/S \cong M^{\mathfrak{C}}$, $M = T\times S$, and either T is transitive on $\mathcal{P}$ or T induces a G-normal partition of $\mathcal{P}$ with 169 parts of size 49. The latter case can not occur because the corresponding CASE, i.e., CASE 9, has been ruled out in (b). Thus T is transitive on $\mathcal{P}$, then by Lemma 3.8 $\mathrm{Soc}(G^{\mathfrak{C}}) = A_7^2$ or $PSL(2,7)^2$, and $c \mid |T|$. So $\mathrm{Soc}(G^{\mathfrak{C}}) = A_7^2$. As a result, $\mathfrak{C} = \mathfrak{C}_1\times\mathfrak{C}_2$, $\mathfrak{C} = \{C_{ij} \mid 1 \le i \le 7, 1 \le j \le 7\}$, $A_7^2 \le G^{\mathfrak{C}} \le S_7 \wr S_2$ in product action. $\mathrm{Soc}(G^{\mathfrak{C}})$ acts on $\mathfrak{C}$ as follow. Let $(h_1,h_2) \in A_7^2 \le G^{\mathfrak{C}}$, $C_{ij} \in \mathfrak{C}$. Then $C_{ij}^{(h_1,h_2)} = C_{i^{h_1}j^{h_2}}$. Let P be a Sylow 5-subgroup of G. Since $5 \nmid b$, $P \le G_\lambda$ for some $\lambda \in \mathcal{L}$. Moreover, because $5 \nmid |GL(2,13)|$, then if P fixes some C_{ij} setwise, it must fix C_{ij} piontwise. Hence $\mathrm{Fix}_{\mathcal{P}}(P)$ is a union of $\mathfrak{C}$-classes. We assume that $P = \langle(12345)\rangle \times \langle(12345)\rangle$, so $\mathrm{Fix}_{\mathfrak{C}}(P) = \{C_{ij} | i,j = 6 \text{ or } 7\}$. Thus $|\mathrm{Fix}_{\mathfrak{C}}(P)| = 4$. Since the intersection type is $(1^{14}, 2^2, 3^{14})$ in LINE 20 and $(1^{28}, 2^7, 4^7)$ in LINE 21, and $P \le G_\lambda$, similar to the proof of CASE 9, we can rule out these two possibilities.

This completes the proof of Proposition 5.1. □

6 The case where $k^{(r)} = 9$

Proposition 6.1 *There is no non-trivial line-transitive point-imprimitive finite linear space $\mathcal{S}$ with the Fang-Li parameter $k^{(r)}$ is 9.*

Proof. Suppose for the contrary that there are line-transitive point-imprimitive symmetric designs with $k^{(r)}$ is 9. Then we need to deal with 106 CASES listed in Table 2. Here the unstated notations in the last collum of Table 2 please refer to Table 1. We prove Proposition 6.1 in 7 steps.

Step 1. Rule out 66 CASES.

There are 6 CASES satisfing $k \le 12$ and ruled out by [20, Theorem 1.1], 54 CASES ruled out by Lemma 3.1. For each of these 54 CASES, we have $k^2 - k + 1 > r$ and there exists a prime $p > k$ such that $p \mid |G|$, but $p \nmid v(v-1)$ which contracts Lemma 3.1. The prime p will be listed in the last column of Table 2.

For CASES 86, 87, 89, 92, 104 and 105, we have $G^{\mathfrak{C}} \ge A_d$, but $t_{max} = 1$ or 2, a contradiction. The symbol (T, t_{max}) denotes this contradiction.

Step 2. CASES 5, 38, 40, 59, 73, 91, 96 and 106 can not occur.

In CASES 40, 91 and 96, we have $\delta = 2$. By Lemma 3.4, we know that $G^{\mathfrak{C}}$ is 2-transitive. In CASES 5, 38, 59, 73 and 106, $\delta = 1$, then by Lemma 3.3, $G^{\mathfrak{C}}$ is 2-transitive. Therefore, by the classification theorem of 2-transitive permutation groups ([4, Theorem 5.3]), in each CASE we have that $G^{\mathfrak{C}} \geq A_d$. But $t_{max} = 2$, contradicting Lemma 3.2. Therefore these CASES are ruled out. The symbol (δ, t_{max}) in Table 2 refers to this situation.

Step 3. CASES 10, 14, 17, 23, 29, 33, 34, 44, 46, 49, 50, 63 and 64 can not occur.

In CASES 10, 14 and 17, we have $\gamma = 1$, by Lemma 3.3, G^C is 2-transitive. In CASE 46, $\gamma = 2$, and d is not a prime power, by Lemma 3.4, G^C is 2-transitive. In CASE 23, $\gamma = 1$, by Lemma 3.3, G^C is also 2-transitive. So by [4, Theorem 5.3], for every CASE the bottom group $G^C \geq A_c$.

For CASES 10, 14 and 19, we have $k^2 - k + 1 > r$, and a prime $p > k$ which is listed in Table 2, with $p \mid |G^C|$, but $p \nmid v(v-1)$, contradicting Lemma 3.1.

For any other CASE, there is a prime p which is listed in Table 2, such that p divides $|G|$, but does not divides b. Thus, in each CASE, a Sylow p-subgroup P of G will fix some line $\lambda \in \mathcal{L}$. Let $F = \mathrm{Fix}_{\mathcal{P}}(P)$. For the line size k, let k_1 to be the integer such that $0 \leq k_1 < k$ and $k \equiv k_1 \pmod{p}$. In each CASE $k_1 > 2$ and $(v-k) \nmid p$ guarantee that there is at least one point which is not on λ fixed by P. Then by Lemma 3.5, F induces an linear space $\mathcal{S}|_F = (F, \mathcal{L}|_F)$. For this subdesign, the number of the points $v_0 = |F| \leq k + r - p - 1$, the line size $k_0 = |\lambda \cap F| \geq k_1$, since $v_0 - 1 \geq k_0(k_0 - 1) \geq k_1(k_1 - 1)$, then

$$k_1(k_1 - 1) \leq k + r - p - 1.$$

But this inequality does not hold in every CASE, the desired contradiction.

Step 4. CASES 22 and 30 can not occur.

CASE 22: Since $G^C \geq A_{1370}$ or $\mathrm{PSL}(2, 37^2)$. Let $p = 1367$, $P \in Syl_p(G)$. If $G^C \geq A_{1370}$, then as above paragraph, we can get the desired contradiction. So $G^C \geq \mathrm{PSL}(2, 37^2)$. Since $t_{max} = 1$, we have $G^{\mathfrak{C}} \ngeq A_{17798}$. For $d = 17798 = 22 \times 809$, let $p = 809$, then G has an element g of order 809. Suppose that g has at least one fixed point in $\mathfrak{C}$, then it has at least 809 fixed points and q cycles of length p where $q \leq 21$. By [22, Theorm 13.10], $|\mathrm{Fix}_{\mathfrak{C}}(g)| \leq 4q - 4$, a contradiction. So g has no fixed point on $\mathfrak{C}$. Thus, by [17, Therom 1.1, Table 3], there is no such primitive group, a contradiction.

CASE 30: Since $G^C \geq A_{1070}$ or $\mathrm{PSL}(2, 1069)$. Let $p = 1063$, $P \in Syl_p(G)$. If $G^C \geq A_{1070}$, similar to Step 3, we can get the desired contradiction. Thus $G^C \geq \mathrm{PSL}(2, 1069)$. Because the prime $p = 1069 \nmid d$, there is a part $C \in \mathfrak{C}$, such that $P \leq G_C$, so $P^C < G^C \leq \mathrm{PGL}(2, 1069)$. Since $\mathrm{PGL}(2, 1069)$ is sharply 3-transitive, the nontrivial subgroup P^C can fix at most 2 points of C. But according to Lemma 3.6, $|\mathrm{Fix}_C(P)| \geq 3$, a contradiction.

Step 5. CASES 1, 2, 4, 6, 7, 11, 27, 32, 47 and 65 can not occur.

For each CASE, we have that $k > 2x + \frac{3}{2} + \sqrt{4x - \frac{7}{4}}$. If the partition is G-normal, by Lemma 3.7, then c is a prime power, a contradiction. So $G^{\mathfrak{C}} \cong G$ is quasiprimitive and almost simple.

For Cases 1, 2, 4, 7, 11, 32, 47 and 65, since $\mathrm{lcm}(b,v) \mid |G|$, then $G \geq A_d$, but in each Case, $t_{max} = 1$ or 2, contradicting Lemma 3.2. For Case 6, we have $\delta = 1$, by Lemma 3.3, $G^{\mathfrak{C}}$ is 2-transitive. Then by [4], we can find that $G^{\mathfrak{C}} \geq A_{8090}$ or $\mathrm{PSL}(2, 8089)$. Furthermore, $b \mid |G|$, so $G^{\mathfrak{C}} \geq A_{8090}$, as a result, the prime $8087 \mid |G|$, this contradicts Lemma 3.1. ForCase 27, since $t_{max} = 2$, then $\mathrm{Soc}(G^{\mathfrak{C}}) = \mathrm{PSL}(5,5)$, it follows that $G_C = G^{\mathfrak{C}}_C = 5^4.\mathrm{GL}(4,5)$. However, from $\mathrm{Soc}(G^C) = \mathrm{PSL}(4,3), \mathrm{PSU}(4,2)$ or A_{40}, we have $3^3 \mid |G_C|$, this is a contradiction.

Step 6. Cases 35, 51, 56, 69 and 97 can not occur.

(1) The partition of each Case is G-normal. Otherwise, $G^{\mathfrak{C}} \cong G$ is an almost simple primitive group of degree d, then $[v,b] \mid |G^{\mathfrak{C}}|$, there is no such group in Case 35, 56 or 97. Now $G \geq A_{103}$ in Case 51, but here t_{max} is 2, a contradiction. So the partition is G-normal relative to K.

(2) Since $x < 8$ and $\mathfrak{C}$ is G-normal in Case 51, it is ruled out by [19, Theorem 1.6]. For Case 69, we have that $k > 2x + \frac{3}{2} + \sqrt{4x - \frac{7}{4}}$. By Lemma 3.7, $K = Z_{103}$ is semiregular, and $Y \cap K = K = S, G/Y \leq \mathrm{Aut}(S) = Z_{102}$. Since $52 \mid |G|$, but $52 \nmid 102$, thus $52 \mid |Y|$, hence $Y^{\mathfrak{C}} \neq 1$. Therefore, by Lemma 3.8, there exists a subgroup M of G, such that $S < M \leq Y, M/S \cong M^{\mathfrak{C}}$, and $M = T \times S$. By [13, Table B.4], $\mathrm{Soc}(G^{\mathfrak{C}}) = \mathrm{PSL}(3,3).2$ or A_{52}. Also we have that $(c,d) = 1$ and by [18, Theorem 5.1.4], the Schur multiplier of $\mathrm{PSL}(3,3)$, A_{52} has no section isomorphic to S, so either T is transitive on $\mathcal{P}$ or T induces a new partition of $\mathcal{P}$ with 103 parts of size 52. The latter case can not occur because there is no corresponding Case with $(d,c) = (52,103)$ in Table 2. Thus T is transitive on $\mathcal{P}$, and $v \mid |T|$, a contradiction. Cases 35 and 56 can be ruled out by the similar method.

For Case 97, if K is not semiegular, by [19, Corollary 4.2] and [19, Lemma 5.1], we have that

$$1 + \frac{r}{k} d_1(d_1 - 1) \leq |\mathrm{Fix}_{\mathcal{P}}(K_\alpha)| \leq d,$$

a contradiction.Thus K is semiregular. Since $K \trianglelefteq G^C$, then K is the unique soluble minimal regular normal subgroup of primitive group G^C, hence $K = Z_{257}$, and $Y \cap K = K = S, G/Y \leq \mathrm{Aut}(S) = Z_{256}$. Since $13 \mid |G^{\mathfrak{C}}|, 13 \nmid |G/Y|$, so $Y^{\mathfrak{C}} \neq 1$. By Lemma 3.8, there exists a normal subgroup M of G, such that $S < M \leq Y, M/S \cong M^{\mathfrak{C}}$, $M = T \times S$, by [13, Tble B.4], $\mathrm{Soc}(G^{\mathfrak{C}}) = A_{65}$, $\mathrm{PSL}(2, 2^6)$, $\mathrm{PSL}(2, 5^2)$, $\mathrm{PSU}(3, 2^2)$ or $Sz(2^3)$, the condition of Lemma 3.8 holds by [18, Theorem 5.1.4], therefore either T is transitive on $\mathcal{P}$ or T induces a G-normal partition of $\mathcal{P}$ with 257 parts of size 40. The latter case can not occur because there is no corresponding Case. Thus T is transitive on $\mathcal{P}$, and $v \mid |T|$, which is impossible.

Step 7. Cases 85 and 101 can not occur.

Case 101: Since $\delta = 1$, then $G^{\mathfrak{C}}$ is 2-transitive, hence $\mathrm{Soc}(G^{\mathfrak{C}}) = \mathrm{PSL}(4,3)$ or A_{40}. If $G \cong G^{\mathfrak{C}}$, then $71 || G^{\mathfrak{C}}|$, a contradiction. So the partition is G-normal. Suppose K is semiregular, then c must be a prime power, this contradiction leads to that K is not semiregular. By [19, Prop. 4.3],

$$d_1 \leq \frac{1}{2} + \sqrt{2k - \frac{7}{4}}.$$

So in this Case the intersection type must be $(2^{10}, 3^{10}, 4^{10})$, i.e., $d_1 = 0$. Now by [19, Prop. 5.2(c)], Case 101 is ruled out.

Case 85: It can be ruled out by the similar method of Case 101.

This complete the proof of Proposition 6.1. ☐

Corollary 6.2 *Let $\mathcal{S}$ be a non-trivial finite linear space with the Fang-Li parameter $k^{(r)}$ is 9. If $G^{\mathfrak{C}} \leq Aut(\mathcal{S})$ is line-transitive, then G is point-primitive.*

Proof of Theorem 1.1 This follows from Propositions 5.1 and 6.1. ☐

References

[1] A. Betten, A. Delandtsheer, M. Law, A. C. Niemeyer, C. E. Praeger, and S. L. Zhou, Linear spaces with a line-transitive point-imprimitive automorphism group and fang-li parameter $\gcd(k, r)$ at most eight. (*available at http://arxiv.org/abs/math.CO/0701629).*

[2] A. Betten, A. Delandtsheer, M. Law, A. C. Niemeyer, C. E. Praeger, and S. L. Zhou, Finite line-transitive linear spaces: theory and search strategies, *Acta Mathematica Sinica, English Series*, **25**(9): 1399-1436, 2009.

[3] A. R. Block, On the orbits of collineation groups, *Math. Z.*, **96**: 33-49, 1967.

[4] P. J. Cameron. Finite permutation groups and finite simple groups. *Bull. London Math. Soc.*, **13(1)**: 1-22, 1981.

[5] A. R. Camina and S. Mishcke, Line-transitive automorphism groups of linear spaces, *Electron. J. Combin.* **3(1)**: Research Paper 3, 16 pp. (electronic), 1996.

[6] A. R. Camina and J. Siemons, Block transitive automorphism groups of $2-(v, k, 1)$ block designs, *J. Combin. Theory Ser. A,* **51(2):** 268-276, 1989.

[7] A. R. Camina and C. E. Praeger, Line-transitive, point quasiprimitive automorphism groups of finite linear spaces are affine or almost simple, *Aequationes Math.*,**61(3)**:221-232, 2001.

[8] H. J. Coutts, M. Quick, and C. M. Roney-Dougal, The primitive permutation groups of degree less than 4096, *Comm. Algebra*, **39(10)**, 3526-3546, 2011.

[9] G. Cresp, Serching for line-transitive, point-imprimitive linear spaces, Honours Dissertation, The Uninversity of Westen Australia, Perth, 2001. (*available at http://arxiv.org/abs/math.CO/0604532).*

[10] A. Delandtsheer, Line-primitive automorphism groups of finite linear spaces, *European J. Combin.*, **10(2)**: 161-169, 1989.

[11] A. Delandtsheer and J. Doyen, Most block-transitive t-designs are point-primitive, *Geom. Dedicata*, 29(3):307-310, 1989.

[12] A. Delandtsheer, A. C. Niemeyer, and C. E. Praeger, Finite line-transitive linear spaces: parameters and normal point-partitions, *Adv. Geom.*, **3(4)**: 469-485, 2003.

[13] J. D. Dixon and B. Mortimer, Permutation groups, Graduate Texts in Mathematics, No.163, *Springer-Verlag*, New York, 1996.

[14] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4*; 2005, `(\protect\vrule width0pt\protect\href{http://www.gap-system.org}{http://www.gap-syst`

[15] W. D. Fang and H. L. Li, A generalization of the Camina-Gagen theorem, *J. Math. (Wuhan)***13(4)**: 437-442, 1993. (in Chinese).

[16] H. L. Li and W.J. Liu, Line-primitive 2-$(v, k, 1)$ desgins with $k/(k, v) \leq 10$, *J. Combin. Theroy Ser. A,* **93,** 153-167, 2001.

[17] M. W. Liebeck and J. Saxl, Primitive permutation groups containing an element of large prime order, *J. London Math. Soc.* **(2), 31(2):**237-249, 1985.

[18] P. Kleidman and M. Liebeck, The subgroup structure of the finite classical groups, London Mathematical Society Lecture Note Series, vol.129, *Cambiridge University Press*, Combridge, 1990.

[19] C. E. Praeger and N. D. Tuan, Inequalities involving the Delandtsheer-Doyen parameters for finite line-transitive linear spaces, *J. Combin. Theory Ser.A*, **102(1)**: 268-276, 2003.

[20] C. E. Praeger and S. L. Zhou, Classification of line-transitive point-imprimitive linear spaces with line size at most 12, *Designs, Codes and Crypt.*, **47**: 99-111, 2008.

[21] C. M. Roney-Donyal, The primitive permutation groups of degree less than 2500, *J Algebra*, **292**: 154-183, 2000.

[22] H. Wielandt, Finite permutation groups, Academic Press, New York, 1964.

7 Appendix

Table 1: Potential parameter sequences for $k^{(r)} = 10$

Line	Case	$d \cdot c$	(x, y)	(γ, δ)	$k^{(v)} \cdot k^{(r)}$	$b^{(v)} \cdot b^{(r)}$	int type	$t_{\max}$	Reference
1	1	$7 \cdot 13$	$(6, 3)$	$(12, 6)$	$1 \cdot 10$	$91 \cdot 1$	$(1, 2^3, 3)$	1	PG$(2, 9)$, [20]
2		$7 \cdot 13$	$(6, 3)$	$(12, 6)$	$1 \cdot 10$	$91 \cdot 1$	$(1^4, 3^2)$	1	
3		$7 \cdot 13$	$(6, 3)$	$(12, 6)$	$1 \cdot 10$	$91 \cdot 1$	$(1^6, 4)$	7	
4	2	$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(3^6, 6^2)$	2	(B, 127)
5		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(2^9, 3, 9)$	2	
6		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^3, 4^3, 5^3)$	2	
7		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^3, 2^3, 5^3, 6)$	2	
8		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^3, 2^6, 3^2, 9)$	2	
9		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^6, 4^3, 6^2)$	2	
10		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^6, 3, 5^3, 6)$	2	
11		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^6, 2^3, 6^3)$	2	
12		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^6, 2^3, 3^3, 9)$	2	
13		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^9, 3, 6^3)$	2	
14		$9 \cdot 129$	$(48, 3)$	$(32, 2)$	$3 \cdot 10$	$387 \cdot 4$	$(1^9, 3^4, 9)$	2	
15	3	$11 \cdot 41$	$(4, 1)$	$(8, 2)$	$1 \cdot 10$	$451 \cdot 5$	$(1^2, 2^4)$	2	non-existence, [9]
16		$11 \cdot 41$	$(4, 1)$	$(8, 2)$	$1 \cdot 10$	$451 \cdot 5$	$(1^5, 2, 3)$	2	
17	4	$13 \cdot 7$	$(3, 6)$	$(6, 12)$	$10 \cdot 1$	$91 \cdot 1$	$(1^4, 2^3)$	1	PG$(2, 9)$, [20]
18		$13 \cdot 7$	$(3, 6)$	$(6, 12)$	$10 \cdot 1$	$91 \cdot 1$	$(1^7, 3)$	1	
19	5	$41 \cdot 11$	$(1, 4)$	$(2, 8)$	$1 \cdot 10$	$451 \cdot 5$	$(1^8, 2)$	2	non-existence, [9]
20	6	$49 \cdot 169$	$(49, 14)$	$(14, 4)$	$7 \cdot 10$	$1183 \cdot 12$	$(1^{14}, 2^7, 3^{14})$	1	KSXY
21		$49 \cdot 169$	$(49, 14)$	$(14, 4)$	$7 \cdot 10$	$1183 \cdot 12$	$(1^{28}, 2^7, 4^7)$	1	
22	7	$111 \cdot 441$	$(36, 9)$	$(8, 2)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{18}, 2^{36})$	2	(T, 109)
23		$111 \cdot 441$	$(36, 9)$	$(8, 2)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{27}, 2^{27}, 3^3)$	2	
24		$111 \cdot 441$	$(36, 9)$	$(8, 2)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{36}, 2^{18}, 3^6)$	2	
25		$111 \cdot 441$	$(36, 9)$	$(8, 2)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{45}, 2^9, 3^9)$	2	
26		$111 \cdot 441$	$(36, 9)$	$(8, 2)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{54}, 3^{12})$	2	
27		$111 \cdot 441$	$(36, 9)$	$(8, 2)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{81}, 9^1)$	2	
28	8	$129 \cdot 9$	$(3, 48)$	$(2, 32)$	$3 \cdot 10$	$387 \cdot 4$	$(1^{24}, 2^3)$	1	(T, 127)
29		$129 \cdot 9$	$(3, 48)$	$(2, 32)$	$3 \cdot 10$	$387 \cdot 4$	$(1^{27}, 3)$	1	
30	9	$169 \cdot 49$	$(14, 49)$	$(4, 14)$	$7 \cdot 10$	$1183 \cdot 12$	$(1^{42}, 2^{14})$	2	KSXY
31	10	$441 \cdot 111$	$(9, 36)$	$(2, 8)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{72}, 2^9)$	2	(B, 109)
32		$441 \cdot 111$	$(9, 36)$	$(2, 8)$	$9 \cdot 10$	$5439 \cdot 55$	$(1^{81}, 3^3)$	1	
33	11	$1937 \cdot 243$	$(39, 312)$	$(2, 16)$	$39 \cdot 10$	$12069 \cdot 121$	$(1^{312}, 9^{39})$	1	(T, 1933)

Table 2: Potential parameter sequences for $k^{(r)} = 9$

LINE	CASE	$d \cdot c$	(x, y)	(γ, δ)	$k^{(v)} \cdot k^{(r)}$	$b^{(v)} \cdot b^{(r)}$	int type	$t_{\max}$	Reference
1	1	$152 \cdot 152$	$(1, 1)$	$(1, 1)$	$2 \cdot 9$	$11552 \cdot 151$	$(1^{16}, 2)$	2	Step 5
2	2	$314 \cdot 314$	$(2, 2)$	$(1, 1)$	$4 \cdot 9$	$24649 \cdot 313$	$(1^{32}, 2^2)$	2	Step 5
3	3	$800 \cdot 800$	$(5, 5)$	$(1, 1)$	$10 \cdot 9$	$64000 \cdot 799$	$(1^{80}, 2^5)$	2	(T,797)
4		$800 \cdot 800$	$(5, 5)$	$(1, 1)$	$10 \cdot 9$	$64000 \cdot 799$	$(1^{83}, 2^2, 3)$	2	
5	4	$1610 \cdot 1610$	$(10, 10)$	$(1, 1)$	$20 \cdot 9$	$129605 \cdot 1609$	$(1^{160}, 2^{10})$	2	Step 5
6		$1610 \cdot 1610$	$(10, 10)$	$(1, 1)$	$20 \cdot 9$	$129605 \cdot 1609$	$(1^{166}, 2^4, 3^2)$	2	
7	5	$4040 \cdot 4040$	$(25, 25)$	$(1, 1)$	$50 \cdot 9$	$326432 \cdot 4039$	$(1^{400}, 2^{25})$	2	$(\delta,\ t_{max})$
8		$4040 \cdot 4040$	$(25, 25)$	$(1, 1)$	$50 \cdot 9$	$326432 \cdot 4039$	$(1^{415}, 2^{10}, 3^5)$	2	
9	6	$8090 \cdot 8090$	$(50, 50)$	$(1, 1)$	$100 \cdot 9$	$654481 \cdot 8089$	$(1^{800}, 2^{50})$	2	8087
10		$8090 \cdot 8090$	$(50, 50)$	$(1, 1)$	$100 \cdot 9$	$654481 \cdot 8089$	$(1^{830}, 2^{20}, 3^{10})$	2	
11	7	$151 \cdot 76$	$(1, 2)$	$(1, 2)$	$2 \cdot 9$	$5738 \cdot 75$	$(1^{16}, 2)$	2	Step 5
12	8	$799 \cdot 400$	$(5, 10)$	$(1, 2)$	$10 \cdot 9$	$31960 \cdot 399$	$(1^{80}, 2^5)$	2	(T, 797)
13		$799 \cdot 400$	$(5, 10)$	$(1, 2)$	$10 \cdot 9$	$31960 \cdot 399$	$(1^{83}, 2^2, 3)$	2	
14	9	$1771 \cdot 886$	$(11, 22)$	$(1, 2)$	$22 \cdot 9$	$71323 \cdot 885$	$(1^{176}, 2^{11})$	2	(B,883)
15	10	$8899 \cdot 4450$	$(55, 110)$	$(1, 2)$	$110 \cdot 9$	$360005 \cdot 4449$	$(1^{880}, 2^{55})$	2	(B,4447)
16		$8899 \cdot 4450$	$(55, 110)$	$(1, 2)$	$110 \cdot 9$	$360005 \cdot 4449$	$(1^{913}, 2^{22}, 3^{11})$	2	
17	11	$149 \cdot 38$	$(1, 4)$	$(1, 4)$	$2 \cdot 9$	$2831 \cdot 37$	$(1^{16}, 2)$	2	Step 5
18	12	$797 \cdot 200$	$(5, 20)$	$(1, 4)$	$10 \cdot 9$	$15940 \cdot 199$	$(1^{80}, 2^5)$	2	199
19		$797 \cdot 200$	$(5, 20)$	$(1, 4)$	$10 \cdot 9$	$15940 \cdot 199$	$(1^{83}, 2^2, 3)$	2	
20	13	$2093 \cdot 524$	$(13, 52)$	$(1, 4)$	$26 \cdot 9$	$42182 \cdot 523$	$(1^{208}, 2^{13})$	2	(B,523)
21	14	$10517 \cdot 2630$	$(65, 260)$	$(1, 4)$	$130 \cdot 9$	$212767 \cdot 2629$	$(1^{1040}, 2^{65})$	2	2621
22	15	$796 \cdot 160$	$(5, 25)$	$(1, 5)$	$10 \cdot 9$	$12736 \cdot 159$	$(1^{80}, 2^5)$	2	(B, 157)
23	16	$1604 \cdot 230$	$(10, 70)$	$(1, 7)$	$20 \cdot 9$	$18446 \cdot 229$	$(1^{160}, 2^{10})$	1	(T, 1601)
24	17	$12944 \cdot 1850$	$(80, 560)$	$(1, 7)$	$160 \cdot 9$	$149665 \cdot 1849$	$(1^{1280}, 2^{80})$	1	1847
25	18	$793 \cdot 100$	$(5, 40)$	$(1, 8)$	$10 \cdot 9$	$7930 \cdot 99$	$(1^{80}, 2^5)$	2	(T,787)
26	19	$13753 \cdot 1720$	$(85, 680)$	$(1, 8)$	$170 \cdot 9$	$139148 \cdot 1719$	$(1^{1360}, 2^{85})$	2	(B,1709)
27	20	$791 \cdot 80$	$(5, 50)$	$(1, 10)$	$10 \cdot 9$	$6328 \cdot 79$	$(1^{80}, 2^5)$	2	(T, 787)
28	21	$628 \cdot 58$	$(4, 44)$	$(1, 11)$	$8 \cdot 9$	$4553 \cdot 57$	$(1^{64}, 2^4)$	1	(T,619)
29	22	$17798 \cdot 1370$	$(110, 1430)$	$(1, 13)$	$220 \cdot 9$	$110833 \cdot 1369$	$(1^{1760}, 2^{110})$	1	1367,809
30	23	$18607 \cdot 1330$	$(115, 1610)$	$(1, 14)$	$230 \cdot 9$	$107597 \cdot 1329$	$(1^{1840}, 2^{115})$	2	1327
31	24	$785 \cdot 50$	$(5, 80)$	$(1, 16)$	$10 \cdot 9$	$3925 \cdot 49$	$(1^{80}, 2^5)$	1	(T, 787)
32	25	$134 \cdot 8$	$(1, 19)$	$(1, 19)$	$2 \cdot 9$	$536 \cdot 7$	$(1^{16}, 2)$	1	(T, 131)
33	26	$3212 \cdot 170$	$(20, 380)$	$(1, 19)$	$40 \cdot 9$	$13651 \cdot 169$	$(1^{320}, 2^{20})$	1	(T, 3209)
34	27	$781 \cdot 40$	$(5, 100)$	$(1, 20)$	$10 \cdot 9$	$3124 \cdot 39$	$(1^{80}, 2^5)$	2	3^3
35	28	$1588 \cdot 70$	$(10, 230)$	$(1, 23)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{160}, 2^{10})$	1	(T, 1583)
36	29	$28315 \cdot 1090$	$(175, 4550)$	$(1, 26)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{2800}, 2^{175})$	1	1087
37	30	$29933 \cdot 1070$	$(185, 5180)$	$(1, 28)$	$370 \cdot 9$	$86563 \cdot 1069$	$(1^{2960}, 2^{185})$	1	1063
38	31	$115 \cdot 4$	$(1, 38)$	$(1, 38)$	$2 \cdot 9$	$230 \cdot 3$	$(1^{16}, 2)$	1	(T, 113)
39	32	$761 \cdot 20$	$(5, 200)$	$(1, 40)$	$10 \cdot 9$	$1522 \cdot 19$	$(1^{80}, 2^5)$	2	Step 5
40	33	$49349 \cdot 950$	$(305, 15860)$	$(1, 52)$	$610 \cdot 9$	$76855 \cdot 949$	$(1^{4880}, 2^{305})$	1	947
41	34	$52585 \cdot 940$	$(325, 18200)$	$(1, 56)$	$650 \cdot 9$	$76046 \cdot 939$	$(1^{5200}, 2^{325})$	1	937
42	35	$76 \cdot 151$	$(2, 1)$	$(2, 1)$	$2 \cdot 9$	$5738 \cdot 75$	$(1^{14}, 2^2)$	2	KSXY
43	36	$400 \cdot 799$	$(10, 5)$	$(2, 1)$	$10 \cdot 9$	$31960 \cdot 399$	$(1^{70}, 2^{10})$	2	(B, 797)
44	37	$886 \cdot 1771$	$(22, 11)$	$(2, 1)$	$22 \cdot 9$	$71323 \cdot 885$	$(1^{154}, 2^{22})$	2	(T, 883)
45	38	$4450 \cdot 8899$	$(110, 55)$	$(2, 1)$	$110 \cdot 9$	$360005 \cdot 4449$	$(1^{770}, 2^{110})$	2	$(\delta,\ t_{max})$
46	39	$35 \cdot 35$	$(1, 1)$	$(2, 2)$	$1 \cdot 9$	$1225 \cdot 17$	$(1^7, 2)$	2	$k < 12$
47	40	$4895 \cdot 4895$	$(121, 121)$	$(2, 2)$	$121 \cdot 9$	$198025 \cdot 2447$	$(1^{847}, 2^{121})$	2	(δ, t_{max})
48	41	$5785 \cdot 2893$	$(143, 286)$	$(2, 4)$	$143 \cdot 9$	$117035 \cdot 1446$	$(1^{1001}, 2^{143})$	1	(B,2887)
49	42	$7120 \cdot 2035$	$(176, 616)$	$(2, 7)$	$176 \cdot 9$	$82325 \cdot 1017$	$(1^{1232}, 2^{176})$	2	(B,2029)
50	43	$31 \cdot 7$	$(1, 5)$	$(2, 10)$	$1 \cdot 9$	$217 \cdot 3$	$(1^7, 2)$	1	$k < 12$
51	44	$9790 \cdot 1507$	$(242, 1573)$	$(2, 13)$	$242 \cdot 9$	$60965 \cdot 753$	$(1^{1694}, 2^{242})$	1	1499
52	45	$29 \cdot 5$	$(1, 7)$	$(2, 14)$	$1 \cdot 9$	$145 \cdot 2$	$(1^7, 2)$	2	$k < 12$
53	46	$10235 \cdot 1463$	$(253, 1771)$	$(2, 14)$	$253 \cdot 9$	$59185 \cdot 731$	$(1^{1771}, 2^{253})$	2	1599
54	47	$433 \cdot 55$	$(11, 88)$	$(2, 16)$	$11 \cdot 9$	$2165 \cdot 27$	$(1^{77}, 2^{11})$	1	Step 5
55	48	$1768 \cdot 187$	$(44, 418)$	$(2, 19)$	$44 \cdot 9$	$7514 \cdot 93$	$(1^{308}, 2^{44})$	1	(T, 1759)
56	49	$15575 \cdot 1199$	$(385, 5005)$	$(2, 26)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{2695}, 2^{385})$	1	1193
57	50	$16465 \cdot 1177$	$(407, 5698)$	$(2, 28)$	$407 \cdot 9$	$47615 \cdot 588$	$(1^{2849}, 2^{407})$	1	1171
58	51	$103 \cdot 52$	$(6, 12)$	$(3, 6)$	$4 \cdot 9$	$1339 \cdot 17$	$(1^{24}, 2^6)$	2	$x < 8$
59	52	$535 \cdot 268$	$(30, 60)$	$(3, 6)$	$20 \cdot 9$	$7169 \cdot 89$	$(1^{120}, 2^{30})$	2	(B, 263)
60	53	$46 \cdot 10$	$(3, 15)$	$(3, 15)$	$2 \cdot 9$	$230 \cdot 3$	$(1^{12}, 2^3)$	2	(T,43)
61	54	$856 \cdot 172$	$(48, 240)$	$(3, 15)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{192}, 2^{48})$	1	(T, 853)
62		$856 \cdot 172$	$(48, 240)$	$(3, 15)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{256}, 4^8)$	1	
63	55	$1177 \cdot 148$	$(66, 528)$	$(3, 24)$	$44 \cdot 9$	$3959 \cdot 49$	$(1^{264}, 2^{66})$	1	(T, 1171)
64		$1177 \cdot 148$	$(66, 528)$	$(3, 24)$	$44 \cdot 9$	$3959 \cdot 49$	$(1^{352}, 4^{11})$	1	
65	56	$38 \cdot 149$	$(4, 1)$	$(4, 1)$	$2 \cdot 9$	$2831 \cdot 37$	$(1^{10}, 2^4)$	2	KSXY
66	57	$200 \cdot 797$	$(20, 5)$	$(4, 1)$	$10 \cdot 9$	$15940 \cdot 199$	$(1^{50}, 2^{20})$	2	(T, 119)
67	58	$524 \cdot 2093$	$(52, 13)$	$(4, 1)$	$26 \cdot 9$	$42182 \cdot 523$	$(1^{130}, 2^{52})$	2	(T, 523)
68	59	$2630 \cdot 10517$	$(260, 65)$	$(4, 1)$	$130 \cdot 9$	$212767 \cdot 2629$	$(1^{650}, 2^{260})$	2	$(\delta,\ t_{max})$
69	60	$2893 \cdot 5785$	$(286, 143)$	$(4, 2)$	$143 \cdot 9$	$117035 \cdot 1446$	$(1^{715}, 2^{286})$	2	(T, 288)
70	61	$17 \cdot 17$	$(2, 2)$	$(4, 4)$	$1 \cdot 9$	$289 \cdot 4$	$(1^5, 2^2)$	1	$k < 12$
71	62	$4208 \cdot 2405$	$(416, 728)$	$(4, 7)$	$208 \cdot 9$	$48655 \cdot 601$	$(1^{1040}, 2^{416})$	1	(B, 2399)
72	63	$5786 \cdot 1781$	$(572, 1859)$	$(4, 13)$	$286 \cdot 9$	$36031 \cdot 445$	$(1^{1430}, 2^{572})$	2	1777
73	64	$6049 \cdot 1729$	$(598, 2093)$	$(4, 14)$	$299 \cdot 9$	$34979 \cdot 432$	$(1^{1495}, 2^{598})$	2	1723
74	65	$257 \cdot 65$	$(26, 104)$	$(4, 16)$	$13 \cdot 9$	$1285 \cdot 16$	$(1^{65}, 2^{26})$	1	Step 5

continued on next page

Line	Case	$d \cdot c$	(x, y)	(γ, δ)	$k^{(v)} \cdot k^{(r)}$	$b^{(v)} \cdot b^{(r)}$	int type	$t_{\max}$	Reference
75	66	$160 \cdot 796$	$(25, 5)$	$(5, 1)$	$10 \cdot 9$	$12736 \cdot 159$	$(1^{40}, 2^{25})$	2	(T, 157)
76	67	$320 \cdot 146$	$(50, 110)$	$(5, 11)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{80}, 2^{50})$	1	(T, 317)
77	68	$352 \cdot 136$	$(55, 143)$	$(5, 13)$	$22 \cdot 9$	$2176 \cdot 27$	$(1^{88}, 2^{55})$	1	(T, 349)
78	69	$52 \cdot 103$	$(12, 6)$	$(6, 3)$	$4 \cdot 9$	$1339 \cdot 17$	$(1^{12}, 2^{12})$	2	KSXY
79		$52 \cdot 103$	$(12, 6)$	$(6, 3)$	$4 \cdot 9$	$1339 \cdot 17$	$(1^{24}, 3^{4})$	2	
80	70	$268 \cdot 535$	$(60, 30)$	$(6, 3)$	$20 \cdot 9$	$7169 \cdot 89$	$(1^{60}, 2^{60})$	2	(T, 263)
81		$268 \cdot 535$	$(60, 30)$	$(6, 3)$	$20 \cdot 9$	$7169 \cdot 89$	$(1^{120}, 3^{20})$	1	
82		$268 \cdot 535$	$(60, 30)$	$(6, 3)$	$20 \cdot 9$	$7169 \cdot 89$	$(1^{120}, 2^{20}, 5^{4})$	1	
83	71	$469 \cdot 235$	$(105, 210)$	$(6, 12)$	$35 \cdot 9$	$3149 \cdot 39$	$(1^{105}, 2^{105})$	1	(T, 467)
84		$469 \cdot 235$	$(105, 210)$	$(6, 12)$	$35 \cdot 9$	$3149 \cdot 39$	$(1^{210}, 3^{35})$	1	
85		$469 \cdot 235$	$(105, 210)$	$(6, 12)$	$35 \cdot 9$	$3149 \cdot 39$	$(1^{210}, 2^{35}, 5^{7})$	1	
86	72	$230 \cdot 1604$	$(70, 10)$	$(7, 1)$	$20 \cdot 9$	$18446 \cdot 229$	$(1^{40}, 2^{70})$	2	B,1601)
87		$230 \cdot 1604$	$(70, 10)$	$(7, 1)$	$20 \cdot 9$	$18446 \cdot 229$	$(1^{80}, 2^{40}, 4^{5})$	2	
88		$230 \cdot 1604$	$(70, 10)$	$(7, 1)$	$20 \cdot 9$	$18446 \cdot 229$	$(1^{100}, 2^{10}, 3^{20})$	2	
89		$230 \cdot 1604$	$(70, 10)$	$(7, 1)$	$20 \cdot 9$	$18446 \cdot 229$	$(1^{120}, 2^{10}, 4^{10})$	2	
90	73	$1850 \cdot 12944$	$(560, 80)$	$(7, 1)$	$160 \cdot 9$	$149665 \cdot 1849$	$(1^{320}, 2^{560})$	2	$(\delta,\ t_{max})$
91		$1850 \cdot 12944$	$(560, 80)$	$(7, 1)$	$160 \cdot 9$	$149665 \cdot 1849$	$(1^{640}, 2^{320}, 4^{40})$	2	
92		$1850 \cdot 12944$	$(560, 80)$	$(7, 1)$	$160 \cdot 9$	$149665 \cdot 1849$	$(1^{800}, 2^{80}, 3^{160})$	2	
93		$1850 \cdot 12944$	$(560, 80)$	$(7, 1)$	$160 \cdot 9$	$149665 \cdot 1849$	$(1^{960}, 2^{80}, 4^{80})$	2	
94		$1850 \cdot 12944$	$(560, 80)$	$(7, 1)$	$160 \cdot 9$	$149665 \cdot 1849$	$(1^{1280}, 8^{20})$	2	
95	74	$2035 \cdot 7120$	$(616, 176)$	$(7, 2)$	$176 \cdot 9$	$82325 \cdot 1017$	$(1^{352}, 2^{616})$	2	(T, 2029)
96		$2035 \cdot 7120$	$(616, 176)$	$(7, 2)$	$176 \cdot 9$	$82325 \cdot 1017$	$(1^{704}, 2^{352}, 4^{44})$	2	
97		$2035 \cdot 7120$	$(616, 176)$	$(7, 2)$	$176 \cdot 9$	$82325 \cdot 1017$	$(1^{880}, 2^{88}, 3^{176})$	2	
98		$2035 \cdot 7120$	$(616, 176)$	$(7, 2)$	$176 \cdot 9$	$82325 \cdot 1017$	$(1^{1056}, 2^{88}, 4^{88})$	2	
99		$2035 \cdot 7120$	$(616, 176)$	$(7, 2)$	$176 \cdot 9$	$82325 \cdot 1017$	$(1^{1408}, 8^{22})$	2	
100	75	$2405 \cdot 4208$	$(728, 416)$	$(7, 4)$	$208 \cdot 9$	$48655 \cdot 601$	$(1^{416}, 2^{728})$	2	(T, 2399)
101		$2405 \cdot 4208$	$(728, 416)$	$(7, 4)$	$208 \cdot 9$	$48655 \cdot 601$	$(1^{1040}, 2^{104}, 3^{208})$	2	
103		$2405 \cdot 4208$	$(728, 416)$	$(7, 4)$	$208 \cdot 9$	$48655 \cdot 601$	$(1^{1248}, 2^{104}, 4^{104})$	2	
104		$2405 \cdot 4208$	$(728, 416)$	$(7, 4)$	$208 \cdot 9$	$48655 \cdot 601$	$(1^{1664}, 8^{26})$	1	
105	76	$3145 \cdot 2752$	$(952, 1088)$	$(7, 8)$	$272 \cdot 9$	$31820 \cdot 393$	$(1^{544}, 2^{952})$	2	(T, 3137)
106		$3145 \cdot 2752$	$(952, 1088)$	$(7, 8)$	$272 \cdot 9$	$31820 \cdot 393$	$(1^{1088}, 2^{544}, 4^{68})$	2	
107		$3145 \cdot 2752$	$(952, 1088)$	$(7, 8)$	$272 \cdot 9$	$31820 \cdot 393$	$(1^{1360}, 2^{136}, 3^{272})$	2	
108		$3145 \cdot 2752$	$(952, 1088)$	$(7, 8)$	$272 \cdot 9$	$31820 \cdot 393$	$(1^{1632}, 2^{136}, 4^{136})$	2	
109		$3145 \cdot 2752$	$(952, 1088)$	$(7, 8)$	$272 \cdot 9$	$31820 \cdot 393$	$(1^{2176}, 8^{34})$	2	
110	77	$100 \cdot 793$	$(40, 5)$	$(8, 1)$	$10 \cdot 9$	$7930 \cdot 99$	$(1^{10}, 2^{40})$	2	(B,787)
111	$100 \cdot 793$	$(40, 5)$	$(8, 1)$	$10 \cdot 9$	$7930 \cdot 99$	$(1^{40}, 2^{10}, 3^{10})$	2		
112	78	$1720 \cdot 13753$	$(680, 85)$	$(8, 1)$	$170 \cdot 9$	$139148 \cdot 1719$	$(1^{170}, 2^{680})$	2	(T, 1709)
113		$1720 \cdot 13753$	$(680, 85)$	$(8, 1)$	$170 \cdot 9$	$139148 \cdot 1719$	$(1^{680}, 2^{170}, 3^{170})$	2	
114	79	$2752 \cdot 3145$	$(1088, 952)$	$(8, 7)$	$272 \cdot 9$	$31820 \cdot 393$	$(1^{272}, 2^{1088})$	1	(B, 3137)
115		$2752 \cdot 3145$	$(1088, 952)$	$(8, 7)$	$272 \cdot 9$	$31820 \cdot 393$	$(1^{1088}, 2^{272}, 3^{272})$	1	
116	80	$80 \cdot 791$	$(50, 5)$	$(10, 1)$	$10 \cdot 9$	$6328 \cdot 79$	$(1^{20}, 2^{20}, 3^{10})$	2	(B, 787)
117	81	$7 \cdot 31$	$(5, 1)$	$(10, 2)$	$1 \cdot 9$	$217 \cdot 3$	$(1^{2}, 2^{2}, 3)$	2	$k < 12$
118	82	$58 \cdot 628$	$(44, 4)$	$(11, 1)$	$8 \cdot 9$	$4553 \cdot 57$	$(2^{32}, 4^{2})$	2	(B, 619)
119		$58 \cdot 628$	$(44, 4)$	$(11, 1)$	$8 \cdot 9$	$4553 \cdot 57$	$(1^{8}, 2^{20}, 3^{8})$	2	
120		$58 \cdot 628$	$(44, 4)$	$(11, 1)$	$8 \cdot 9$	$4553 \cdot 57$	$(1^{16}, 2^{20}, 4^{4})$	2	
121		$58 \cdot 628$	$(44, 4)$	$(11, 1)$	$8 \cdot 9$	$4553 \cdot 57$	$(1^{24}, 2^{8}, 3^{8}, 4^{2})$	2	
122		$58 \cdot 628$	$(44, 4)$	$(11, 1)$	$8 \cdot 9$	$4553 \cdot 57$	$(1^{32}, 2^{8}, 4^{6})$	2	
123	83	$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(2^{80}, 4^{5})$	2	(B,317)
124		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{20}, 2^{50}, 3^{20})$	2	
125		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{20}, 2^{70}, 5^{4})$	1	
126		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{40}, 2^{50}, 4^{10})$	2	
127		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{60}, 2^{20}, 3^{20}, 4^{5})$	2	
128		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{60}, 2^{40}, 4^{5}, 5^{4})$	1	
129		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{80}, 2^{10}, 3^{20}, 5^{4})$	1	
130		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{80}, 2^{20}, 4^{15})$	2	
131		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{80}, 2^{30}, 5^{8})$	1	
132		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{100}, 2^{10}, 4^{10}, 5^{4})$	1	
133		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{120}, 4^{5}, 5^{8})$	1	
134		$146 \cdot 320$	$(110, 50)$	$(11, 5)$	$20 \cdot 9$	$2336 \cdot 29$	$(1^{120}, 2^{20}, 10^{2})$	1	
135	84	$235 \cdot 469$	$(210, 105)$	$(12, 6)$	$35 \cdot 9$	$3149 \cdot 39$	$(2^{105}, 3^{35})$	1	(B, 467)
136		$235 \cdot 469$	$(210, 105)$	$(12, 6)$	$35 \cdot 9$	$3149 \cdot 39$	$(1^{70}, 2^{105}, 7^{5})$	1	
137		$235 \cdot 469$	$(210, 105)$	$(12, 6)$	$35 \cdot 9$	$3149 \cdot 39$	$(1^{105}, 3^{70})$	2	
138		$235 \cdot 469$	$(210, 105)$	$(12, 6)$	$35 \cdot 9$	$3149 \cdot 39$	$(1^{175}, 4^{35})$	1	
139		$235 \cdot 469$	$(210, 105)$	$(12, 6)$	$35 \cdot 9$	$3149 \cdot 39$	$(1^{175}, 3^{35}, 7^{5})$	1	
140	85	$1370 \cdot 17798$	$(1430, 110)$	$(13, 1)$	$220 \cdot 9$	$110833 \cdot 1369$	$(2^{770}, 4^{110})$	2	STEP 7
141		$1370 \cdot 17798$	$(1430, 110)$	$(13, 1)$	$220 \cdot 9$	$110833 \cdot 1369$	$(1^{440}, 2^{110}, 3^{440})$	2	
142		$1370 \cdot 17798$	$(1430, 110)$	$(13, 1)$	$220 \cdot 9$	$110833 \cdot 1369$	$(1^{660}, 2^{110}, 3^{220}, 4^{110})$	2	
143		$1370 \cdot 17798$	$(1430, 110)$	$(13, 1)$	$220 \cdot 9$	$110833 \cdot 1369$	$(1^{880}, 2^{110}, 4^{220})$	2	
144		$1370 \cdot 17798$	$(1430, 110)$	$(13, 1)$	$220 \cdot 9$	$110833 \cdot 1369$	$(1^{1100}, 2^{330}, 11^{20})$	2	
145	86	$1507 \cdot 9790$	$(1573, 242)$	$(13, 2)$	$242 \cdot 9$	$60965 \cdot 753$	$(2^{847}, 4^{121})$	2	(T, t_{max})
146		$1507 \cdot 9790$	$(1573, 242)$	$(13, 2)$	$242 \cdot 9$	$60965 \cdot 753$	$(1^{484}, 2^{121}, 3^{484})$	2	
147		$1507 \cdot 9790$	$(1573, 242)$	$(13, 2)$	$242 \cdot 9$	$60965 \cdot 753$	$(1^{726}, 2^{121}, 3^{242}, 4^{121})$	2	
148		$1507 \cdot 9790$	$(1573, 242)$	$(13, 2)$	$242 \cdot 9$	$60965 \cdot 753$	$(1^{968}, 2^{121}, 4^{242})$	2	
149		$1507 \cdot 9790$	$(1573, 242)$	$(13, 2)$	$242 \cdot 9$	$60965 \cdot 753$	$(1^{1210}, 2^{363}, 11^{22})$	2	
150	87	$1781 \cdot 5786$	$(1859, 572)$	$(13, 4)$	$286 \cdot 9$	$36031 \cdot 445$	$(2^{1001}, 4^{143})$	1	(T, t_{max})

continued on next page

Line	Case	$d \cdot c$	(x, y)	(γ, δ)	$k^{(v)} \cdot k^{(r)}$	$b^{(v)} \cdot b^{(r)}$	int type	$t_{\max}$	Reference
151		$1781 \cdot 5786$	$(1859, 572)$	$(13, 4)$	$286 \cdot 9$	$36031 \cdot 445$	$(1^{572}, 2^{143}, 3^{572})$	1	
152		$1781 \cdot 5786$	$(1859, 572)$	$(13, 4)$	$286 \cdot 9$	$36031 \cdot 445$	$(1^{858}, 2^{143}, 3^{286}, 4^{143})$	1	
153		$1781 \cdot 5786$	$(1859, 572)$	$(13, 4)$	$286 \cdot 9$	$36031 \cdot 445$	$(1^{1144}, 2^{143}, 4^{286})$	1	
154		$1781 \cdot 5786$	$(1859, 572)$	$(13, 4)$	$286 \cdot 9$	$36031 \cdot 445$	$(1^{1430}, 2^{429}, 11^{26})$	1	
155	88	$136 \cdot 352$	$(143, 55)$	$(13, 5)$	$22 \cdot 9$	$2176 \cdot 27$	$(2^{77}, 4^{11})$	1	(B,349)
156		$136 \cdot 352$	$(143, 55)$	$(13, 5)$	$22 \cdot 9$	$2176 \cdot 27$	$(1^{44}, 2^{11}, 3^{44})$	1	
157		$136 \cdot 352$	$(143, 55)$	$(13, 5)$	$22 \cdot 9$	$2176 \cdot 27$	$(1^{66}, 2^{11}, 3^{22}, 4^{11})$	1	
158		$136 \cdot 352$	$(143, 55)$	$(13, 5)$	$22 \cdot 9$	$2176 \cdot 27$	$(1^{88}, 2^{11}, 4^{22})$	1	
159		$136 \cdot 352$	$(143, 55)$	$(13, 5)$	$22 \cdot 9$	$2176 \cdot 27$	$(1^{110}, 2^{33}, 11^{2})$	1	
160	89	$1330 \cdot 18607$	$(1610, 115)$	$(14, 1)$	$230 \cdot 9$	$107597 \cdot 1329$	$(1^{230}, 2^{230}, 3^{460})$	2	(T, t_{max})
161		$1330 \cdot 18607$	$(1610, 115)$	$(14, 1)$	$230 \cdot 9$	$107597 \cdot 1329$	$(1^{690}, 2^{230}, 4^{230})$	2	
162	90	$5 \cdot 29$	$(7, 1)$	$(14, 2)$	$1 \cdot 9$	$145 \cdot 2$	$(1, 2, 3^{2})$	2	$k < 12$
163		$5 \cdot 29$	$(7, 1)$	$(14, 2)$	$1 \cdot 9$	$145 \cdot 2$	$(1^{3}, 2, 4)$	2	
164	91	$1463 \cdot 10235$	$(1771, 253)$	$(14, 2)$	$253 \cdot 9$	$59185 \cdot 731$	$(1^{253}, 2^{253}, 3^{506})$	2	(δ, t_{max})
165		$1463 \cdot 10235$	$(1771, 253)$	$(14, 2)$	$253 \cdot 9$	$59185 \cdot 731$	$(1^{759}, 2^{253}, 4^{253})$	2	
166	92	$1729 \cdot 6049$	$(2093, 598)$	$(14, 4)$	$299 \cdot 9$	$34979 \cdot 432$	$(1^{299}, 2^{299}, 3^{598})$	1	(T, t_{max})
167		$1729 \cdot 6049$	$(2093, 598)$	$(14, 4)$	$299 \cdot 9$	$34979 \cdot 432$	$(1^{897}, 2^{299}, 4^{299})$	1	
168	93	$10 \cdot 46$	$(15, 3)$	$(15, 3)$	$2 \cdot 9$	$230 \cdot 3$	$(2^{3}, 3^{4})$	2	(B, 43)
169		$10 \cdot 46$	$(15, 3)$	$(15, 3)$	$2 \cdot 9$	$230 \cdot 3$	$(1^{2}, 2^{3}, 3^{2}, 4)$	1	
170		$10 \cdot 46$	$(15, 3)$	$(15, 3)$	$2 \cdot 9$	$230 \cdot 3$	$(1^{4}, 2^{3}, 4^{2})$	1	
171	94	$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(2^{48}, 3^{64})$	2	(B, 853)
172		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(2^{96}, 4^{24})$	2	
173		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(2^{128}, 8^{4})$	1	
174		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{32}, 2^{48}, 3^{32}, 4^{16})$	1	
175		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{64}, 3^{64}, 4^{8})$	1	
176		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{64}, 2^{48}, 4^{32})$	1	
177		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{64}, 2^{80}, 4^{8}, 8^{4})$	1	
178		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{96}, 3^{32}, 4^{24})$	1	
179		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{96}, 2^{32}, 3^{32}, 8^{4})$	1	
180		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{128}, 4^{40})$	1	
181		$172 \cdot 856$	$(240, 48)$	$(15, 3)$	$32 \cdot 9$	$4601 \cdot 57$	$(1^{128}, 2^{32}, 4^{16}, 8^{4})$	1	
182	95	$50 \cdot 785$	$(80, 5)$	$(16, 1)$	$10 \cdot 9$	$3925 \cdot 49$	$(1^{10}, 2^{20}, 4^{10})$	2	(B, 773)
183		$50 \cdot 785$	$(80, 5)$	$(16, 1)$	$10 \cdot 9$	$3925 \cdot 49$	$(1^{20}, 3^{20}, 5^{2})$	2	
184		$50 \cdot 785$	$(80, 5)$	$(16, 1)$	$10 \cdot 9$	$3925 \cdot 49$	$(1^{20}, 2^{10}, 3^{10}, 5^{4})$	2	
185		$50 \cdot 785$	$(80, 5)$	$(16, 1)$	$10 \cdot 9$	$3925 \cdot 49$	$(1^{20}, 2^{20}, 5^{6})$	2	
186		$50 \cdot 785$	$(80, 5)$	$(16, 1)$	$10 \cdot 9$	$3925 \cdot 49$	$(1^{40}, 4^{10}, 5^{2})$	2	
187		$50 \cdot 785$	$(80, 5)$	$(16, 1)$	$10 \cdot 9$	$3925 \cdot 49$	$(1^{50}, 5^{8})$	2	
188	96	$55 \cdot 433$	$(88, 11)$	$(16, 2)$	$11 \cdot 9$	$2165 \cdot 27$	$(1^{11}, 2^{22}, 4^{11})$	2	(δ, t_{max})
189	97	$65 \cdot 257$	$(104, 26)$	$(16, 4)$	$13 \cdot 9$	$1285 \cdot 16$	$(1^{13}, 2^{26}, 4^{13})$	1	KSXY
190	98	$8 \cdot 134$	$(19, 1)$	$(19, 1)$	$2 \cdot 9$	$536 \cdot 7$	$(2, 3^{4}, 4)$	3	(B, 131)
191		$8 \cdot 134$	$(19, 1)$	$(19, 1)$	$2 \cdot 9$	$536 \cdot 7$	$(1^{2}, 2, 3^{2}, 4^{2})$	3	
192		$8 \cdot 134$	$(19, 1)$	$(19, 1)$	$2 \cdot 9$	$536 \cdot 7$	$(1^{4}, 2, 4^{3})$	3	
193		$8 \cdot 134$	$(19, 1)$	$(19, 1)$	$2 \cdot 9$	$536 \cdot 7$	$(1^{4}, 2^{4}, 6)$	3	
194	99	$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(2^{20}, 3^{80}, 4^{20})$	2	(B, 3209)
195		$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(2^{80}, 4^{50})$	2	
196		$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(1^{40}, 2^{20}, 3^{40}, 4^{40})$	2	
197		$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(1^{80}, 2^{20}, 4^{60})$	2	
198		$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(1^{80}, 2^{80}, 6^{20})$	2	
199		$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(1^{80}, 2^{100}, 8^{10})$	2	
200		$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(1^{160}, 2^{20}, 4^{10}, 6^{20})$	2	
201		$170 \cdot 3212$	$(380, 20)$	$(19, 1)$	$40 \cdot 9$	$13651 \cdot 169$	$(1^{160}, 2^{40}, 4^{10}, 8^{10})$	2	
202	100	$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(2^{22}, 3^{88}, 4^{22})$	2	(B, 1759)
203		$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(2^{88}, 4^{55})$	2	
204		$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(1^{44}, 2^{22}, 3^{44}, 4^{44})$	2	
205		$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(1^{88}, 2^{22}, 4^{66})$	2	
206		$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(1^{88}, 2^{88}, 6^{22})$	2	
207		$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(1^{88}, 2^{110}, 8^{11})$	2	
208		$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(1^{176}, 2^{22}, 4^{11}, 6^{22})$	2	
209		$187 \cdot 1768$	$(418, 44)$	$(19, 2)$	$44 \cdot 9$	$7514 \cdot 93$	$(1^{176}, 2^{44}, 4^{11}, 8^{11})$	2	
210	101	$40 \cdot 781$	$(100, 5)$	$(20, 1)$	$10 \cdot 9$	$3124 \cdot 39$	$(2^{10}, 3^{10}, 4^{10})$	2	STEP 7
211		$40 \cdot 781$	$(100, 5)$	$(20, 1)$	$10 \cdot 9$	$3124 \cdot 39$	$(1^{40}, 5^{10})$	2	
212	102	$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(2^{20}, 4^{35})$	2	(B, 1583)
213		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(2^{50}, 4^{5}, 6^{10})$	2	
214		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(2^{60}, 4^{5}, 8^{5})$	2	
215		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{20}, 2^{20}, 3^{20}, 6^{10})$	2	
216		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{20}, 2^{30}, 5^{20})$	2	
217		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{20}, 2^{30}, 3^{20}, 8^{5})$	2	
218		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{40}, 2^{20}, 4^{10}, 6^{10})$	2	
219		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{40}, 2^{30}, 4^{10}, 8^{5})$	2	
220		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{60}, 4^{5}, 5^{20})$	2	
221		$70 \cdot 1588$	$(230, 10)$	$(23, 1)$	$20 \cdot 9$	$5558 \cdot 69$	$(1^{60}, 3^{20}, 4^{5}, 8^{5})$	2	
222	103	$148 \cdot 1177$	$(528, 66)$	$(24, 3)$	$44 \cdot 9$	$3959 \cdot 49$	$(2^{44}, 3^{88}, 11^{4})$	1	(B, 1171)
223		$148 \cdot 1177$	$(528, 66)$	$(24, 3)$	$44 \cdot 9$	$3959 \cdot 49$	$(2^{88}, 5^{44})$	1	
224		$148 \cdot 1177$	$(528, 66)$	$(24, 3)$	$44 \cdot 9$	$3959 \cdot 49$	$(1^{44}, 4^{88})$	1	
225		$148 \cdot 1177$	$(528, 66)$	$(24, 3)$	$44 \cdot 9$	$3959 \cdot 49$	$(1^{88}, 2^{44}, 4^{44}, 11^{4})$	1	

continued on next page

Line	Case	$d \cdot c$	(x, y)	(γ, δ)	$k^{(v)} \cdot k^{(r)}$	$b^{(v)} \cdot b^{(r)}$	int type	$t_{\max}$	Reference
226		$148 \cdot 1177$	$(528, 66)$	$(24, 3)$	$44 \cdot 9$	$3959 \cdot 49$	$(1^{132}, 2^{88}, 11^{8})$	1	
227	104	$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(3^{350}, 4^{350}, 5^{140})$	2	$(\mathrm{T},\ t_{max})$
228		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(3^{700}, 5^{140}, 7^{50})$	2	
229		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(2^{350}, 4^{350}, 5^{210})$	2	
230		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(2^{350}, 3^{350}, 5^{210}, 7^{50})$	2	
231		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(2^{700}, 5^{280}, 7^{50})$	2	
232		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{350}, 3^{350}, 5^{350})$	2	
233		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{350}, 2^{350}, 5^{420})$	2	
234		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{350}, 2^{350}, 4^{350}, 7^{100})$	2	
235		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{350}, 2^{350}, 3^{350}, 7^{150})$	2	
236		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{350}, 2^{700}, 5^{70}, 7^{150})$	2	
237		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{700}, 4^{350}, 5^{140}, 7^{50})$	2	
238		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{700}, 3^{350}, 5^{140}, 7^{100})$	2	
239		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{700}, 2^{350}, 5^{210}, 7^{100})$	2	
240		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{700}, 2^{350}, 3^{350}, 10^{70})$	2	
241		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{700}, 2^{700}, 5^{70}, 10^{70})$	2	
242		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{1050}, 5^{350}, 7^{50})$	2	
243		$1090 \cdot 28315$	$(4550, 175)$	$(26, 1)$	$350 \cdot 9$	$88181 \cdot 1089$	$(1^{1050}, 2^{350}, 7^{200})$	2	
244	105	$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(3^{385}, 4^{385}, 5^{154})$	2	$(T,\ t_{max})$
245		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(3^{770}, 5^{154}, 7^{55})$	2	
246		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(2^{385}, 4^{385}, 5^{231})$	2	
247		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(2^{385}, 3^{385}, 5^{231}, 7^{55})$	2	
248		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(2^{770}, 5^{308}, 7^{55})$	2	
249		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{385}, 3^{385}, 5^{385})$	2	
250		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{385}, 2^{385}, 5^{462})$	2	
251		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{385}, 2^{385}, 4^{385}, 7^{110})$	2	
252		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{385}, 2^{385}, 3^{385}, 7^{165})$	2	
253		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{385}, 2^{770}, 5^{77}, 7^{165})$	2	
254		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{770}, 4^{385}, 5^{154}, 7^{55})$	2	
255		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{770}, 3^{385}, 5^{154}, 7^{110})$	2	
256		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{770}, 2^{385}, 5^{231}, 7^{110})$	2	
257		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{770}, 2^{385}, 3^{385}, 10^{77})$	2	
258		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{770}, 2^{770}, 5^{77}, 10^{77})$	2	
259		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{1155}, 5^{385}, 7^{55})$	2	
260		$1199 \cdot 15575$	$(5005, 385)$	$(26, 2)$	$385 \cdot 9$	$48505 \cdot 599$	$(1^{1155}, 2^{385}, 7^{220})$	2	
261	106	$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(5^{650}, 10^{260})$	2	$(\delta,\ t_{max})$
262		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(4^{650}, 5^{260}, 13^{150})$	2	
263		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(3^{650}, 5^{260}, 10^{130}, 13^{100})$	2	
264		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(2^{650}, 5^{390}, 10^{130}, 13^{100})$	2	
265		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(2^{650}, 4^{650}, 15^{130})$	2	
266		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(2^{650}, 3^{650}, 13^{200})$	2	
267		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(1^{650}, 8^{650})$	2	
268		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(1^{650}, 5^{650}, 13^{150})$	2	
269		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(1^{650}, 3^{650}, 5^{260}, 15^{130})$	2	
270		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(1^{650}, 2^{650}, 10^{390})$	2	
271		$940 \cdot 52585$	$(18200, 325)$	$(56, 1)$	$650 \cdot 9$	$76046 \cdot 939$	$(1^{650}, 2^{650}, 5^{390}, 15^{130})$	2	